

## Quiz for Wave Medical RN Student

**1) Tetrodotoxin is caused by the consumption of which of the following?**

- a) Puffer Fish
- b) Fugu
- c) Blow Fish
- d) All of the above
- e) None of the above

**2) In what section would you find immediate, life saving protocol?**

- a) Table of Context
- b) Home Index
- c) Acute Care Reference
- d) Quick Reference
- e) None of the above

**3) Out of the drugs listed, which combination should never be used because of its serious, potential life threatening interaction?**

- a) Zomig and Bromocriptine
- b) Zomig and Viagra
- c) Zomig and Kepivance
- d) All of the above
- e) None of the above

**4) Out of the drugs listed, which combination should never be used because of its serious, potential life threatening interaction?**

- a) Rifampin and St. Johns Wort
- b) Rifampin and Allegra-D
- c) Rifampin and Voriconazole
- d) All of the above
- e) None of the above

---

**5) What is the peak flow of a 6 year old girl who is 4' tall?**

- a) 187 L/min
- b) 200 L/min
- c) 214 L/min
- d) 227 L/min
- e) None of the above

**6) What is the Total Body Water of a 150 lbs adult female?**

- a) 34.01 L
- b) 46.83 L
- c) 50.82 L
- d) 50.83 L
- e) None of the above

**7) What is the adult dosing for Vitamin B1 in males?**

- a) 1.1 mg/d
- b) 1.2 mg/d
- c) .3 g/d
- d) .3 mg/d
- e) None of the Above

**8) What does the abbreviation DC stand for in medical terminology?**

- a) Discharge from wound
- b) Discharge Patient
- c) Ductal Carcinoma in Situ
- d) Dilated Cardiomyopathy
- e) None of the above

**9) In treating someone with a Subarachnoid Hemorrhage, the nurse should do which of the following?**

- a) Anticipate intubation for major bleeds, increase ICP and loss of protective airway reflexes
- b) Monitor VS and neuro signs closely
- c) Provide comfort measures, including Analgesics, non-narcotics preferred to prevent masking neuro changes, and a cool cloth over eyes
- d) All of the above
- e) None of the above

**10) To maintain a constant 0.025 mg/hr infusion of Buprenex, using 15 mcg of the drug in a 2 ml bag, set the IV drip rate to which of the following?**

- a) 200 ml/hr
- b) 205 ml/hr
- c) 210 ml/hr
- d) 215 ml/hr
- e) None of the above

**11) The reportable range of C-Reactive Protein in humans is which of the following?**

- a) 0.01-20 mg/dL
- b) 0.2-10 mg/dL
- c) 0.3-20 mg/dL
- d) 0.5-30 mg/dL
- e) None of the above

**12) What is the normal reference range of Hemoglobin in a pregnant female?**

- a) > 11 g/dL
- b) 12-16 g/dL
- c) 14-18 g/dL
- d) > 18 g/dL
- e) None of the above

---

13) Which of the following is NOT a stage of Whooping Cough?

- a) Catarrhal Stage
- b) Paroxysmal Stage
- c) Expectorant Stage
- d) Convalescent stage
- e) None of the above

14) To initiate a 0.05-0.1 mcg/kg/min IV infusion of Norepinephrine in a 55 lb, 8 year old boy, using 4 mg of the drug in a 1000 ml bag, set the drip rate to which of the following?

- a) 35 ml/hr
- b) 45 ml/hr
- c) 51.25 ml/hr
- d) 60 ml/hr
- e) None of the above

15) Which of the following drug combinations should never be used because of its serious, potential life threatening interaction?

- a) Parnate and Penbutolol
- b) Parnate and Taro Carbamazepine
- c) Parnate and Phenobarbital
- d) All of the Above
- e) None of the Above

## Answers

- 1) d
- 2) c
- 3) b
- 4) c
- 5) c
- 6) a
- 7) b
- 8) b
- 9) d
- 10) a
- 11) c
- 12) a
- 13) c
- 14) e
- 15) e